



ТОЧНЫЕ

В XVII-XVIII

ВЕКАХ

ВЫПОЛНИЛИ СТУДЕНТКИ 1
КУРСА СВ-13

ДАВЫДОВА ЕЛИЗАВЕТА

МАРЧЕНКО АЛЕКСАНДРА

$$\frac{\partial}{\partial a} \ln J_{a, \sigma^2}(\xi_1) = \frac{(\xi_1 - a)}{\sigma^2} f_{a, \sigma^2}(\xi_1) = \frac{1}{\sqrt{2\pi\sigma}} \exp\left(-\frac{(\xi_1 - a)^2}{2\sigma^2}\right)$$
$$\int_{\mathbb{R}_n} T(x) \cdot \frac{\partial}{\partial \theta} f(x, \theta) dx = M\left(T(\xi) \cdot \frac{\partial}{\partial \theta} \ln L(\xi, \theta)\right)$$
$$\int_{\mathbb{R}_n} T(x) \cdot \left(\frac{\partial}{\partial \theta} \ln L(x, \theta)\right) \cdot f(x, \theta) dx = \int_{\mathbb{R}_n} T(x) \cdot \left(\frac{\partial}{\partial \theta} \frac{f(x, \theta)}{f(x, \theta)}\right) f(x, \theta) dx$$

